

# UKSCA Position Statement: Youth Resistance Training

**Rhodri S. Lloyd**, PhD, CSCS\*D, ASCC<sup>1</sup>; **Avery D. Faigenbaum**, EdD, FACSM, CSCS\*D<sup>2</sup>; **Gregory D. Myer**, PhD, CSCS\*D, FACSM<sup>3</sup>; **Michael H Stone**, PhD, ASCC, FNCSA, FUKSCA<sup>4</sup>; **Jon L. Oliver**, PhD<sup>5</sup>; **Ian Jeffreys**, PhD, CSCS\*D, ASCC, FNCSA<sup>6</sup>; **Jeremy Moody**, PhD, ASCC<sup>5</sup>; **Clive Brewer**, MSc, CSCS, ASCC<sup>7</sup>; **Kyle Pierce** EdD, CSCS<sup>8</sup>

## Preamble

This manuscript will serve as the official position statement of the United Kingdom Strength and Conditioning Association (UKSCA) for youth resistance training. The authorship team for this manuscript were selected from the fields of paediatric exercise science, physical education, elite sport, and sports medicine. Prior to publishing, the manuscript was reviewed in detail and subsequently endorsed by the UKSCA Board of Directors.

## Operational Definitions

Prior to discussing the literature surrounding the potential benefits and concerns associated with youth resistance training, it is pertinent to define key terminologies used throughout the manuscript.

- *Childhood* represents the developmental period of life from the end of infancy to the beginning of adolescence. The term *children* refers to girls and boys (generally up to the age of 11 years and 13 years respectively) who have not developed secondary sex characteristics.
- The term *adolescence* refers to a period of life between childhood and adulthood. Although adolescence is a more difficult period to define in terms of chronological age due to differential maturation rates,<sup>126</sup> girls 12-18 years and boys 14-18 years are generally considered adolescents.
- The terms *youth* and *young athletes* represent global terms which include both children and adolescents
- *Growth* is typically viewed as a quantifiable change in body composition, the size of the body as a whole, or the size of specific regions of the body.<sup>21</sup>
- *Maturation* refers to the highly variable timing and tempo of progressive change within the human body from childhood to adulthood, and which, in addition to growth, influences overall physical performance capabilities.<sup>21</sup>
- *Training age* refers to the number of years an athlete has been involved in a formalised training programme.<sup>119</sup>



*Rhodri S. Lloyd is a senior lecturer in Strength and Conditioning at the University of Gloucestershire. He is a fully accredited strength and conditioning coach with the UKSCA, and holds certified strength and conditioning status with distinction from the NSCA.*

*Rhodri's research interests surround strength and conditioning for young athletes, and the application of long-term athlete development models in the physical preparation of youths. Elected onto the UKSCA Board of Directors in 2011, he is the inaugural convenor of the UKSCA Youth Training Special Interest Group, and is paediatric lead for the association.*

1 Faculty of Applied Sciences, University of Gloucestershire, UK

2 Department of Health and Exercise Science, The College of New Jersey, USA

3 Department of Pediatrics and Orthopaedic Surgery, College of Medicine, University of Cincinnati, USA

4 Center of Excellence for Sport Science and Coach Education, East Tennessee State University, USA

5 Cardiff School of Sport, Cardiff Metropolitan University (UWIC), UK

6 Faculty of Health, Sport and Science, University of Glamorgan, UK

7 Widnes Vikings Rugby League Club, UK

8 Department of Kinesiology and Health Science, Louisiana State University, Shreveport, USA

Address for Correspondence:

Rhodri S. Lloyd, PhD, ASCC, CSCS\*D, rhodri@uksca.org.uk

- Resistance training (also called strength training) refers to a specialised method of conditioning whereby an individual is working against a wide range of resistive loads in order to enhance health, fitness and performance. Forms of resistance training include the use of body weight, weight machines, free weights (barbells and dumbbells), elastic bands and medicine balls.
- *Weightlifting* is a sport, which involves the performance of the snatch and clean and jerk lifts in competition.<sup>183</sup> The sport of weightlifting is distinct from the terms resistance training, strength training or weightlifting training.

## Introduction

Despite outdated misperceptions that resistance training was unsafe or inappropriate for youth,<sup>2</sup> there is now a compelling body of scientific evidence that supports its use by children and adolescents for a wide range of performance, health, and injury reducing benefits.<sup>3,10,17,18,69,167,185</sup> Research has indicated that various forms of resistance training can elicit significant performance improvements in muscular strength,<sup>19</sup> power production,<sup>69</sup> running velocity,<sup>134</sup> change-of-direction speed,<sup>188</sup> and general motor performance,<sup>18</sup> in youth. From a health perspective, evidence suggests that resistance training can make positive alterations in overall body composition,<sup>177</sup> reduce abdominal and trunk fat,<sup>200</sup> improve insulin-sensitivity in overweight adolescents,<sup>179</sup> and enhance cardiac function in obese children.<sup>150</sup> Importantly, it has also been suggested that regular participation in an appropriately designed exercise programme, which includes resistance training, can enhance bone mineral density<sup>13</sup> and reduce sports-related injury risk in young athletes.<sup>143,193</sup> This would appear to be an important consideration given that the European Network for Sports Injury Prevention<sup>15</sup> previously estimated nearly 1.3 million cases of sports injuries requiring hospitalisation in Europe were for children under the age of 15 years. Additionally, muscular fitness and resistance training have been associated with positive psychological health and wellbeing in children and adolescents.<sup>58,103,160,195,206</sup>

Despite the apparent health, fitness and performance benefits for school-age youth, there is little reference to resistance training in the physical education national curriculums within the UK. Of note, no information regarding age-appropriate resistance training is included for primary aged children (11 years and under), and resistance training guidelines for secondary school age children (11 years and upwards) are limited and inconsistent. Compounding this issue, recent evidence indicates that muscular strength levels of children within the United Kingdom are decreasing.<sup>40</sup> Such a tendency of declines in muscular strength levels is commensurate with other European countries.<sup>35,173</sup> Progressive resistance training under the supervision of qualified health and fitness professionals can offer a method for reversing this undesirable trend while encouraging participation in resistance training as an ongoing lifestyle choice. The importance of effective education by qualified personnel is essential, as positive early experiences in physical education have been associated with lifelong physical activity.<sup>111</sup> Therefore, the current manuscript will dispel the myths surrounding youth resistance training, and provide guidance and support for those individuals responsible for the long-term development and general wellbeing of school-age youth.

## The Effects Of Growth And Maturation On The Development Of Muscular Strength During Childhood And Adolescence

It has previously been established that muscular strength development is a multi-dimensional fitness component that is influenced by a combination of muscular, neural and biomechanical factors.<sup>45</sup> Due to the non-linear development of physiological processes, such as stature and body mass during childhood and adolescence, the assessment and monitoring of muscular strength can be a challenging task.<sup>80</sup> Similarly, a non-linear pattern emerges when examining the development of physical performance qualities in younger populations.<sup>21</sup> Assessments of muscular strength in children and adolescents have traditionally been attained from isometric testing, and this data suggest that strength increases in a relatively linear fashion throughout childhood for both boys and girls.<sup>30</sup> As children reach the onset of puberty, they experience rapid growth and increases in muscular strength.<sup>161</sup> During this period, sex-differences in muscular strength levels start to emerge, with boys demonstrating accelerated gains as a result of the adolescent spurt, and girls appearing to continue to develop in a more linear fashion.<sup>21</sup> Factors inherently responsible for increases in strength during childhood appear to be related to the development of the central nervous system. Specifically, improvements in motor unit recruitment, firing frequency, synchronisation, and neural myelination are all deemed to enhance neuromuscular performance.<sup>112,170</sup> Conversely, natural strength gains during adolescence are driven by further neural development, but also structural and architectural changes resulting largely from increased sex androgen concentrations, including testosterone, growth hormone and insulin-like growth factor (IGF-I).<sup>126</sup> Further increases in muscle cross-sectional area, muscle pennation angle, and continued muscle fibre type differentiation will all enable adolescents to express greater levels of force, and partly explains the differences in strength between children, adolescents and adults.<sup>189</sup> Sex-related differences in muscular strength are more evident as children enter adulthood, with males consistently outperforming females.<sup>155</sup> Research has indicated that muscle growth will largely explain the disparity between genders, especially for absolute measures of muscular strength and power.<sup>156,151</sup>

It is essential that physical education teachers, paediatric fitness specialists, health care providers, and strength and conditioning coaches who train children and adolescents, are aware of these basic paediatric scientific principles in order to ensure that exercise prescription is always planned according to the unique demands of the individual, irrespective of chronological age. Owing to the highly individualised nature of growth and maturation, children and adolescents of the same chronological age will vary markedly in biological status (possibly up to 4-5 years difference), and consequently, chronological age is deemed a weak indicator of maturational status.<sup>20</sup> Awareness of the potential variation in biological age amongst children of the same chronological age group is a central tenet of most long-term athlete development programmes in a bid to ensure that children are trained according to their biological status, as opposed to age-group classifications.<sup>8,9,11,27,32,119,153</sup> In addition to chronological and biological age, those responsible for the design and

implementation of youth resistance training programmes must take into consideration the training age of the individual.<sup>119</sup> From a developmental perspective, this becomes critically important when training an adolescent who is approaching adulthood, but has no experience of participating in structured resistance training programmes. Conversely, a 10 year old child with a two-year training age, who can demonstrate proficient technical competency on basic exercises, should not be restricted to introductory training methods typically commensurate with their chronological age.

## Health Benefits Of Resistance Training For Youth

The World Health Organisation now recognises physical inactivity as the fourth leading risk factor for global mortality for non-communicable diseases, and supports participation in a variety of physical activities including those that strengthen muscle and bone.<sup>203</sup> Since contemporary youth are not as active as they should be,<sup>54,84,88</sup> children and adolescents should be encouraged to participate regularly in play, games, sports, and planned exercise in the context of school and community activities. Not only is physical activity essential for normal growth and development, but also youth programmes that enhance fundamental movement skill performance early in life appear to build the foundation for an active lifestyle later in life.<sup>12,122,182</sup> Since muscular strength is an essential component of motor skill performance,<sup>126</sup> developing competence and confidence to perform resistance exercise during the growing years may have important long-term implications for health, fitness and sports performance.

Resistance training can offer unique health benefits to children and adolescents when appropriately prescribed and supervised. Regular participation in a youth resistance training programme can have a favourable influence on musculoskeletal health, body composition, and cardiovascular risk factors.<sup>17,57,115,158,178</sup> Moreover, since physical inactivity is a risk factor for activity-related injuries in children,<sup>26</sup> youth who participate regularly in age-appropriate fitness programmes, which include resistance exercise, may be less likely to suffer an injury.<sup>39</sup> These observations are particularly important for sedentary youth whose musculoskeletal system may be ill prepared for the demands of recreational games and sports practice.<sup>60,143,144</sup> The International Olympic Committee recognises the importance of physical activity and sport for youth, and promotes the early identification of fitness deficits in aspiring young athletes, and the proper prescription of training programmes to address individual limitations.<sup>136</sup> From a public health perspective, it is noteworthy that traditional fears and misinformed concerns that resistance training would damage the developing skeleton, have been replaced by reports indicating that childhood may be the opportune time to build bone mass and enhance bone structure by participating in weight-bearing physical activities.<sup>87,197</sup> Fears that resistance training would injure the growth-plates of youths are not supported by scientific reports, which indicate that the mechanical stress placed on the developing growth plates from resistance exercise, or high strain eliciting sports such as weightlifting, are actually beneficial for bone formation and growth.<sup>42,199</sup> While numerous factors, including genetics and nutritional status, influence skeletal health, regular

participation in impact-loading sports and physical activity programmes, which include multi-joint, moderate to high intensity resistance exercise, can help to optimise bone mineral accrual during childhood and adolescence.<sup>1,13,14,25,48,81,102,152,202</sup> Furthermore, no scientific evidence indicates that resistance training will have an adverse effect on linear growth during childhood or adolescence,<sup>125</sup> or reduce eventual height in adulthood.<sup>75,125</sup>

Given the growing prevalence of overweight and obesity among children and adolescents and the associated health-related concerns, the influence of resistance training on the metabolic health, body composition and injury risk profile of overweight and obese youth has received increased attention.<sup>43,44,129,130,179,186,194</sup> Although aerobic exercise is typically prescribed for overweight or obese youth, excess body weight hinders the performance of physical activities such as jogging, and overweight and obese adolescents are more than twice as likely to be injured in sports and other physical activities compared with non-overweight and non-obese adolescents.<sup>130</sup> Furthermore, overweight and obese youth seem to demonstrate significantly lower motor coordination than normal weight youth.<sup>47,123</sup> While the treatment of overweight and obese youth is complex, participation in a multi-faceted treatment programme, including resistance training, may provide a gateway for this population to improve muscle strength, enhance motor coordination and gain confidence in their perceived abilities to be physically active.<sup>181</sup>

The available evidence indicates that resistance training has the potential to offer observable health value to sedentary youth and young athletes, and this type of training can be prescribed to meet the needs of all children and adolescents, regardless of body size or physical ability.

## Injury Prevention Benefits Of Resistance Training For Youth

The total elimination of sport- and physical activity related injuries is an unrealistic goal; however, preparatory resistance training programmes that are appropriately designed and sensibly progressed for youth may help to reduce the likelihood of injuries in these populations. Lehnhard and colleagues<sup>116</sup> were able to significantly reduce injury rates with the addition of a resistance training regimen to a male soccer team. Cahill and Griffith<sup>34</sup> incorporated resistance training into their preseason conditioning for adolescent football teams and reported a reduction in non-serious knee injuries, as well as knee injuries that required surgery, over four competitive seasons. Hejna *et al.*<sup>93</sup> reported that young athletes who incorporated resistance training in their exercise regimen suffered fewer injuries and recovered from injuries with less time spent in rehabilitation as compared to teammates who did not resistance train. Similarly, Soligard *et al.*<sup>180</sup> successfully reduced the risk of severe and overuse injuries in female youth football players, following the implementation of a comprehensive warm-up programme that incorporated resistance-based exercises. Likewise, Emery and Meeuwisse<sup>55</sup> reported a reduction in overall injuries and acute injury incidence in youth football players with the use of an integrative neuromuscular training (INT) programme focused on resistance training. It should be noted that INT refers to a training programme that incorporates general and specific strength and conditioning activities

that improve health- and skill-related physical fitness components.<sup>144</sup>

Resistance training that is focused to address the risk factors associated with youth sport injuries (e.g. low fitness level, muscle imbalances and errors in training) has the potential to reduce overuse injuries by up to 50%.<sup>133,193</sup> For example, several investigations indicate that reduced relative hamstring strength is related to both hamstring strains and ACL injury.<sup>107,145,157,204</sup> Integrative resistance training protocols incorporated into pre-season and in-season conditioning programmes reduced these injury risks, and decreased anterior cruciate ligament injuries in young athletes.<sup>95,96,139,140,141</sup> Even stronger associations of injury reduction are reported for decreased hamstring strains in young adult athletes who utilised hamstring resistance training exercises in preseason conditioning for their respective sports, such as football, rugby, and Australian rules football.<sup>5,6,83,163</sup>

While there is not one combination of resistance training exercises that has proven to optimise training adaptations, it appears that multi-faceted programmes that increase muscle strength, enhance movement mechanics, and improve functional abilities appear to be the most effective strategy for reducing sports-related injuries in young athletes.<sup>49,55,95,137</sup> Clearly, participation in physical activity should not begin with competitive sport but should evolve out of preparatory fitness conditioning that is sensibly progressed over time. Owing to the apparent decline in free time physical activity among children and adolescents,<sup>148,154</sup> it seems that the musculoskeletal system of some aspiring young athletes may not be prepared for the demands of sports practice and competition. Of concern, research has suggested that physical activity levels in youth peak at approximately 6 years of age, and consistently decline throughout childhood and into youth.<sup>192</sup> Consequently, the supporting structures of some young athletes may be ill prepared to handle the demands of weekly sports practice sessions and weekend competitions. Cumulatively, these findings indicate that young athletes should participate regularly in multi-faceted resistance training programmes prior to sports seasons to reduce their risk of sports related injury.

### *Special Resistance Training Considerations for Young Females*

Musculoskeletal growth during puberty, in the absence of corresponding neuromuscular adaptation, may facilitate the development of abnormal joint mechanics and injury risk factors in young girls.<sup>78,97</sup> These intrinsic risk factors, if not addressed at the proper time, may continue through adolescence and into maturity, thus predisposing female athletes to increased risk of injuries.<sup>101,146</sup> In a recent longitudinal study, Ford *et al.*<sup>77</sup> noted that young females who did not participate in resistance training programmes as they matured developed injury risk factors. Conversely, those who did report participation in resistance training activities during maturation were found to have safer movement mechanics and associated increased posterior chain strength.<sup>77</sup>

Well supervised, multifaceted resistance training programmes reduce these abnormal biomechanics<sup>100,138,140,141</sup> and appear to decrease injury rates in female athletes.<sup>95</sup> The findings of a recent meta-analysis revealed that an age-related association between integrative resistance training implementation

and reduction of ACL incidence only occurred in the youngest female athletes.<sup>137</sup> Integrative resistance training utilised to enrich the motor learning environment in early youth may initiate adaptation and help low motor competence children “catch-up” with their peers in these measures.<sup>41,91,171,172</sup> In addition to reduced knee injuries in adolescent<sup>98</sup> and mature female athletes,<sup>147</sup> regular participation in a multi-faceted resistance training programme may also induce measures of the “neuromuscular spurt” which are not typically seen in females.<sup>141</sup> Of potential interest to sports medicine professionals, resistance training timed with growth and development may induce the desired neuromuscular spurt, which may improve sports performance and improve biomechanics related to injury risk in young females.<sup>101,141</sup> Observed relative gains in females may be greater than in males because baseline neuromuscular performance levels are lower on average in females.<sup>79,99,113,141,168</sup>

Although there are many mechanisms to potentially reduce sports related injuries, enhancing muscular fitness and strength using resistance training as a preventative health measure should be considered a cornerstone of year-round, multifaceted training programmes for school age youth.

## Effectiveness Of Youth Resistance Training

As previously documented, children and adolescents will naturally increase muscular strength levels as a result of growth and maturation.<sup>20,21,22,23,126</sup> The appropriate development of strength as a result of training can have important implications for sport and daily life. In order to induce adaptations in muscular strength levels above and beyond those of growth and maturation alone, the magnitude and intensity of training stimulus must be sufficient.<sup>18,50,69,94,126</sup> However, research clearly indicates that appropriately designed, and well supervised resistance training programmes can benefit youth of all ages, with children as young as 5 years of age making noticeable improvements in muscular fitness.<sup>4,65,108,201</sup> Interestingly, whilst the magnitude of absolute strength gains have been reported as being greater in adolescents (ES = 1.91) in comparison to children (ES = 0.81),<sup>19</sup> relative increases in strength appear to be similar across childhood and adolescence.<sup>118,162,165</sup>

The term ‘trainability’ describes the sensitivity of developing athletes to a given training stimulus at varying stages of growth and maturation. Growth and maturation complicate the concept of trainability, as combined they can quite often imitate potential training effects.<sup>16,127,149</sup> Recently, Keiner *et al.*<sup>110</sup> examined the trainability of 141 youth football players (11 to 19 years old). Participants were divided into either a football training only group, or a combined football training and resistance training group. The data indicated a high level of trainability for all age groups (under 19’s, under 17’s, under 15’s and under 13’s). Using absolute and relative back squat measures, two years of resistance training resulted in differences between control and resistance trained groups ranging from 125% in the younger group to 54% in the older group. Interestingly, the strength profiles of the football players was compared against data of 105 youth weightlifters, and it emerged that the young weightlifters were stronger than the soccer players at each age group. Considering the effect of increased strength on reducing injuries<sup>55</sup> and increasing sport

related performance variables such as sprinting, jumping, and general sport performance,<sup>18,31,37,92,117</sup> the investigators suggested that with 4-5 years of training, relative strength levels (1RM kg/kg body mass) in the back squat should be a minimum of 2.0 for late adolescents (16-19 yrs); 1.5 for adolescents (13-15 yrs); and 0.7 for children (11-12 yrs). These findings, in addition to several reviews of the literature<sup>57,59,60,75,119,143,167,185</sup> including three meta-analyses,<sup>19,76,162</sup> highlight the effectiveness of resistance training for sports performance for all youth, and underline that resistance training programmes should not be limited to adult populations.

Despite the growing body of evidence demonstrating that resistance training can lead to established improvements in performance through increases in strength, speed, power and other related characteristics (e.g. rate of force development), an aspect that has caused debate amongst researchers, deals with the degree of hypertrophy that is possible as a result of strength training among children prior to puberty.<sup>24,82,127,131</sup> Existing research suggests that increases in muscular strength are a result of both architectural (muscle size, movement arm length) and neural (voluntary activation level) adaptations.<sup>29,155</sup> However, the mechanisms appear to differ according to the stage of development. The primary mechanisms underlying training-induced gains in muscular strength and related characteristics before puberty appear to depend upon neural adaptations.<sup>126,198</sup> However, among early and particularly late adolescence, the effects of resistance training appear to be a result of additional gains in lean body mass and muscle cross-sectional area, with further alterations in neural mechanisms appearing to be the same as those adaptations experienced by adults.<sup>19,159</sup> Thus, it is among children that very special care should be taken in their development, and coaches should avoid using training methods aimed at increasing overall muscle mass, instead focusing on strength development and movement skill competency.<sup>119</sup> Indeed, among children, from a conceptual standpoint, the adoption of a long-term approach to athletic development should be implemented.<sup>119,121</sup>

Combined, the existing literature highlights a number of important concepts. Firstly, at all ages, appropriate resistance training can clearly result in an increased level of strength and related characteristics.<sup>18,19,51</sup> Within the literature, gains in maximum strength have ranged from approximately 10% to 70%<sup>66</sup> depending upon several factors including the type of training programme, duration of the training programme, and the type of exercise used to measure strength (e.g. isometric versus isoinertial, large muscle mass versus small muscle mass, and partial versus complete range of motion). However, in general, expected strength gains of 30-40% are typically observed in untrained youth following participation in an introductory resistance training programme. Secondly, strength training results in only a minor sex-associated effect on both absolute and relative strength gains among pre-pubertal children, however, the magnitude of effect does appear to be a function of sex in older groups.<sup>118</sup> Thirdly, evidence indicates that more effective programmes lasted more than 8 weeks, involved multiple sets and that generally, gains increased with the frequency of training sessions per week.<sup>18</sup> However, it should be noted that youth with extensive resistance training experience need to follow periodised training

programmes with systematic variation in intensity, volume, frequency and repetition velocity to facilitate continued adaptation. This is supported with evidence from adult based data, that suggests a reduction in training frequency accompanied by an increase in training intensity produces superior results.<sup>164</sup>

### *Weightlifting: an Effective Training Tool for Young Athletes*

Previous literature highlights that weightlifting is considerably safer for children and adolescents than many have generally believed, providing qualified supervision and instruction are available and progression is based on the technical performance of each lift.<sup>33,90,121,142,166,167</sup> Therefore, despite previous misconceptions surrounding its safety, the use of weightlifting as part of a multifaceted youth-based resistance training programme is supported, on the proviso that training prescription is designed in line with the maturational status, training age, and level of motor competency of the young athlete, and that those supervising possess the relevant expertise.

Weightlifting injury rate is reportedly lower than other forms of resistance training and sports in general.<sup>90</sup> For example, Pierce *et al.*<sup>166</sup> observed 70 female and male children participating in weightlifting ranging in age from 7 to 16 years, who regularly completed maximal and near-maximal lifts in competition and training. Over a 12 month period, no days of training were lost as a result of injury from weightlifting,<sup>166</sup> and both boys and girls increased physical strength as measured by weightlifting performance.<sup>166</sup> A more detailed follow-up study of 3 females (13.7 ± 1.2 y) and 8 males (12.5 ± 1.6 y) across a year's competition (534 competition lifts) showed similar results.<sup>33</sup> Both boys and girls showed marked enhancement of weightlifting performance, and no injuries requiring medical attention or loss of training time occurred.<sup>33</sup> Furthermore, these two observational studies indicated that if training and competition are appropriate for the age group, and are properly supervised and sensibly progressed, then weightlifting exercises may provide a unique stimulus for enhancing strength, explosiveness and power performance in school-age youth. Due to the high skill level required to perform weightlifting movements, it is important that individuals responsible for teaching these complex movements to youths, hold the requisite coaching qualifications, and have suitable experience of teaching youths.<sup>121</sup>

Weightlifting training uses a variety of large multi-muscle, multi-joint exercises, such as squats, snatches, and clean and jerks (plus variations of these movements).<sup>121</sup> Weightlifting has been used to examine potential effects of strength-power training on a number of performance and physiological variables among children and adolescents.<sup>33,53,89</sup> A series of studies carried out in the Soviet Union examined the effects of weightlifting training on the physical development, physiology and performance of children and adolescents. The data gleaned from these studies indicate that from 12 years to 22 years of age, weightlifting training can produce positive alterations in body composition, cardiorespiratory variables, various motor fitness parameters (e.g. jumping and sprinting), and overall weightlifting performance.<sup>51,53</sup>

Considering the observational, correlation and longitudinal data as a whole, it is accepted that resistance training (inclusive of weightlifting training) can improve strength and strength related variables

among children and adolescents. These studies also offer evidence as to the efficacy of strength gain alone in producing positive alterations in skill related variables among children and adolescents.<sup>18</sup>

## Resistance Training Guidelines For Children And Adolescents

Whilst recent evidence has indicated that both children and adolescents can make significant gains in muscular strength and other health- and skill-related fitness components,<sup>18,19</sup> it is imperative that suitably qualified personnel are responsible for coaching and instruction of youth-based resistance training programmes.<sup>69,119,196</sup> Such qualifications would include a relevant strength and conditioning qualification (for example, the Accredited Strength and Conditioning Coach (ASCC) status), a strong understanding of paediatric exercise science, and crucially, relevant pedagogic experience and a communication skill-set to enable effective engagement with youth of varying abilities and personalities.<sup>69,119,121</sup>

### Training Variable Considerations

#### Exercise selection

Whilst a range of exercises, performed on a variety of equipment can be prescribed to both children and adolescents, it is vital that the fundamentals of technical competency are prioritised at all times. Researchers have previously highlighted the importance of equipment modifications and habituation when testing youth for any fitness component, and promote the use of 'child-sized' or 'child friendly' machinery wherever possible.<sup>45,46,184</sup> The principles of equipment suitability and familiarity for paediatric testing, should also apply for youths participating in resistance training. Some of the resistance modes available to those prescribing youth resistance training programmes include: bodyweight, machine weights, free weights, resistance bands, medicine balls, and manual resistance,<sup>62</sup> all of which have been successfully implemented within the training programmes of both children and adolescents.<sup>38,51,59,64,65,67,71,73,86,105,120,132,141,174,179,187,188,190,191,209</sup> The selection of resistance modality will largely depend upon the technical ability and baseline fitness levels of the individual. For example, it may be necessary to gradually introduce sedentary youth to simple resistance training exercises via the use of machine-based resistance, in order to enhance their self-perception of competency, before progressing to more complex, free weight based activities. It should be recognised however, that when competency is sufficiently developed, free weight resistance should be used wherever possible, as machine-based resistance has been reported to stimulate lower muscle activation in lower,<sup>176</sup> upper,<sup>175</sup> and whole-body<sup>106</sup> exercises compared to free-weight resistance. Recent evidence demonstrated that whilst a 10-week weight-machine based training programme was successful in increasing isokinetic peak torque of the knee extensors and knee flexors in prepubertal children, it was unsuccessful in improving more dynamic athletic qualities, including vertical jump height and postural sway.<sup>85</sup> Such dynamic qualities can be enhanced in youth with multi-joint, velocity-specific training in the form of free weight resistance training,<sup>71</sup> weightlifting,<sup>166</sup> and dynamic exercise using body weight, such as plyometrics.<sup>120</sup> For youths with a low training age, coaches should employ a range of exercises which are designed to promote the development of muscular strength and

enhance overall fundamental movement competency. Childhood is deemed to be a crucial time in which to develop movement competency, as it is during these formative years that neuromuscular coordination is most susceptible to change.<sup>28</sup> During this stage of development, children will experience rapid brain maturation,<sup>169</sup> and exposing children to key athletic movement patterns at a time where natural strengthening of existing synaptic pathways<sup>36</sup> and synaptic pruning<sup>35</sup> takes place, is considered crucial for long-term athletic development<sup>119</sup> and life-long physical activity.<sup>124</sup> Once the child can demonstrate appropriate technical competency, they should be introduced to more advanced exercises that challenge the child in terms of coordination and require greater levels of force generation. In the case of weightlifting exercises, which by their nature are complex, multi-joint exercises, authors have previously suggested that early exposure should focus on technical development using body weight, modified equipment and light external loads.<sup>69,121,167</sup>

#### Training Volume and Intensity

Volume and intensity are key resistance training variables that are routinely manipulated within a training session, or overall phase of training, depending on the primary training goal of the individual. Volume refers to the total number of times an exercise is performed within a training session, and is comprised of the number of sets, and number of repetitions within each set.<sup>7</sup> Volume can also relate to the overall loading within short- and long-term training blocks, but for the purposes of the current manuscript the former definition will be used. Intensity most commonly refers to the magnitude of resistance that is required to be overcome during a repetition.<sup>7</sup> The relationship between volume and intensity is inverse in nature; the greater the load (intensity), the lower the number of repetitions that can be completed (volume) by the individual.<sup>7</sup> Both variables must be considered synergistically when prescribing resistance training for youth, however, training intensity is arguably more important owing to the injury risk associated with exposing a child or adolescent to excessive external loading at the expense of correct technique.

In order to prescribe appropriate training intensity, coaches typically stipulate a percentage of an athlete's one repetition-maximum (1RM). Research indicates that maximal strength and power testing of children<sup>63</sup> and adolescents<sup>70</sup> is safe and reliable when standardised protocols are implemented and monitored by qualified instructors. Whilst 1RM measurements are routinely used within a research environment or elite level sport, owing to time and equipment constraints, physical education teachers will typically benefit more from the use of predictive equations that estimate 1RM values from sub-maximal loads in youth populations.<sup>104,114,128</sup> However, it should be acknowledged that methods of predicting 1RM values from higher repetition ranges possess less accuracy, in particular when repetition ranges exceed more than 10.<sup>63</sup> Additionally, it should be noted that a child or adolescent must be able to demonstrate sound technical competency irrespective of the RM range selected.

#### Progression of Volume and Intensity

When biologically immature children, or youth with a low training age, first begin to participate in formalised resistance training in schools and recreation

programmes, the use of 1RM measurements (actual or predicted) to determine training intensities will typically be unnecessary. Consequently, teachers and coaches should prescribe an appropriate repetition range to develop technical competency and acquire a base level of adaptation, and over time increase external load until technique is deemed to have sufficiently improved. Therefore, a beginner may be prescribed 1-2 sets of 8-12 repetitions with a light or moderate load (50-70% 1RM or equivalent), however, with increased exposure to formalised resistance training, they may increase prescription to 2-4 sets of 6-10 repetitions with a heavier load (70-85% 1RM). Once technical expertise is sufficiently high, adolescents may be introduced to lower volumes (2-5 sets of 3-5 repetitions) and higher external loads (>85% 1RM). However, it is important to note that not all exercises need to be performed for the same number of sets and repetitions within a training session. For example, a relatively experienced adolescent may begin a resistance training session performing 3 sets of 5 repetitions of a power-oriented exercise (e.g. power clean); then complete 3 sets of 6-8 repetitions of a large compound, multi-joint movement (e.g. back squat); and then finish with 2 sets of 6-10 repetitions of a unilateral exercise (e.g. barbell step-up). Depending on the learning environment, coaches will need to provide feedback to ensure technical competency does not decrease towards the end of a prescribed set. The frequency and mode of feedback will depend to a large degree on the number of individuals training, mode of exercise being performed, and the stage of learning and personality traits of the young athletes involved. For example, when coaching a novice weightlifter, feedback may be provided after individual repetitions;<sup>121</sup> however, for a primary school physical education class of 25 pupils, where the focus of the lesson is aimed at fundamental movement skill development, feedback may be provided less frequently.

### *Rest Intervals During Training Sessions*

While limited evidence exists examining the optimal rest periods for youth-based resistance training, available research indicates that children can recover more quickly from fatigue-inducing resistance exercise.<sup>72,208</sup> It is suggested that children are less likely to suffer muscle damage following such exercise, owing to the increased pliability of their muscle tissue.<sup>56</sup> Rest periods of approximately 1 minute should suffice for inexperienced children, however, this should be increased as children enter adolescence and become more experienced (2-5 minutes), especially if the exercises require high levels of skill, force or power production. Whilst children can recover more quickly from short, intermittent, high intensity training than adults,<sup>72,74,208</sup> as a caveat, teachers and coaches should monitor within-session resistance training performance to ensure correct technique is maintained.

### *Training Frequency*

Training frequency refers to the number of sessions performed within a week. Previous research has indicated that 2-3 sessions per week on non-consecutive days is most appropriate in order to develop muscular strength levels in children and adolescents.<sup>69</sup> Behringer and colleagues<sup>19</sup> recently substantiated these previous recommendations, indicating that across 42 studies (where mean training frequency was  $2.7 \pm 0.8$  sessions per week), training frequency was significantly correlated with increased

resistance training effect. Training frequency may increase as children go through adolescence and approach adulthood, especially for youths in competitive sport. While promoting sampling and exposure to a variety of physical activity experiences to help promote physical development,<sup>119,124</sup> in the interests of athlete welfare and well-being, coaches, parents and educators should be cognisant of the potential difficulties when children are asked to participate in numerous activities and thus accumulating high exercise volumes. For youth participating in competitive sports, in-season resistance training should be viewed as mandatory to maintain muscular strength levels and reduce injury-risk. Physical education classes may offer a suitable time for youths involved in sport to receive such training (providing suitably qualified personnel are responsible for the delivery), as research demonstrates that exposure to resistance training during exercise lessons or physical education classes does not have an adverse effect on after-school performance in adolescent athletes.<sup>68</sup>

### *Repetition Velocity*

Whilst existing guidelines promote the use of controlled, moderate movement velocities for youths, there is arguably a need to promote the intention to move quickly to develop motor unit recruitment patterns and firing frequencies.<sup>205</sup> Variations in repetition velocity may also be indicative of the training age of the child; for example, a child with limited training history may need to perform all exercises with slow to moderate speed to maximise control and ensure correct technical development, whereas an early adolescent with three years training history could be exposed to much greater movement velocities. Repetition velocities may also fluctuate within a session; for example, the movement preparation phase may consist of slow and controlled movements, the main dynamic and explosive exercises (plyometrics or weightlifting derivatives) will involve rapid movement speeds, whilst the main strength development exercises (squatting, deadlifting, pressing and pulling) will typically include slow to moderate movement velocities. However, even when external loads are sufficient to produce slow movement speed, there should arguably be an intention to move explosively wherever possible to promote appropriate neuromuscular adaptations and to maximise the transfer of training effect.<sup>109</sup> Arguably, the development of high velocity movement is especially important for children and adolescents at a time when neural efficiency and motor coordination is most sensitive to change.<sup>35</sup>

*Table 1* provides a summary of suggested guidelines for youth-based resistance training prescription aimed at improving muscular strength levels for a range of abilities and experiences. These are only guidelines, and flexibility will be required in order to ensure prescription of resistance training protocols for youths is at all times individualised. It should be noted that resistance training prescription for youths should always be age-related and not age-determined. Sensible progression should be made from beginner status through to advanced, and those responsible for determining an athlete's level of competency should consider a range of compounding variables, inclusive of maturational stage, training age, fundamental movement skill competency, technical proficiency, existing strength levels and psychosocial factors (such

Table 1. Suggested guidelines for youth-based resistance training prescription

Training Experience	Beginner	Intermediate	Experienced	Advanced
Volume (sets x reps)	1-2 x 8-12	2-4 x 6-10	2-4 x 5-8	2-5 x 2-5
Total number of exercises per session	6-10	3-6	3-6	2-5
Intensity (%1RM)	Body weight, or 50-70% 1RM	60-80%	70-85%	85-100%
Repetition Velocity (speed of movement)	Moderate-Fast	Moderate-Fast	Fast-Maximal	Maximal
Rest intervals (minutes)	1	1-2	2-3	2-5
Frequency (sessions per week)	2-3	2-3	2-4	2-5
Recovery (hours in between sessions)	72-48	72-48	48	48-24

as confidence and self-perception of motor competency). Regardless of training status the level of instruction, education and feedback provided to the child or adolescent will be fundamental to the success of the training programme.

### Summary

A compelling body of scientific evidence supports participation in well-designed youth resistance training programmes that are supervised and instructed by qualified professionals. The current manuscript has added to previous position statements from medical and fitness organisations, and has clearly outlined the performance, health, and injury reducing benefits associated with this training mode for school age youth. In summarising this manuscript, it is the position of the UKSCA that:

1. The use of resistance training by children and adolescents is supported on the proviso that qualified professionals supervise training programmes that are consistent with the needs, goals, and abilities of younger populations.
2. Specifically, the use of weightlifting as a resistance training mode by children and adolescents is supported, providing appropriate equipment and logical progressions are prescribed and implemented by suitably qualified personnel.
3. Parents, teachers and coaches, should recognise the potential health-related benefits of resistance exercise, because youths who do not participate in activities that enhance muscle strength and movement skills, may be at increased risk for negative health outcomes later in life.
4. Well designed resistance training programmes are not only safe for young athletes but may also reduce sports-related injuries.
5. An appropriately designed resistance training programme can elicit noticeable improvements in motor skills, and consequently may positively enhance sports performance.
6. A properly designed resistance training programme can improve and maintain psychological health and wellbeing.
7. Resistance training prescription for children and adolescents should be age-related and not age-determined. Consequently, prescription should be based according to biological status, training age, motor skill competency, technical proficiency, existing strength levels, and psychosocial maturity.

8. Regardless of resistance training mode, training volume and intensity should never be increased at the expense of technical competency.

### References

1. Álvarez-San Emeterio, C., Palacios-Gil Antuñano, N., López-Sobale, A.M. and González-Badillo, J.J. Effect of strength training and the practice of alpine skiing on bone mass density, growth, body composition and the strength and power of the legs of adolescent skiers. *Journal of Strength and Conditioning Research*, 25: 2879–2890. 2011.
2. American Academy of Pediatrics. Weight training and weight lifting: Information for the Pediatrician. *Physician and Sports Medicine*, 11: 157-61. 1983.
3. American College of Sports Medicine. ACSMs Guidelines for Exercise Testing and Prescription. 8th ed. Lippincott Williams and Wilkins, Philadelphia, PA, 2010.
4. Annesi, J., Westcott, W., Faigenbaum, A. and Unruh, J. Effects of a 12 week physical activity program delivered by YMCA after-school counselors (Youth Fit for Life) on fitness and self-efficacy changes in 5–12 year old boys and girls. *Research Quarterly for Exercise and Sport*, 76: 468–476. 2005.
5. Arnason, A., Andersen, T.E., Holme, I., Engebretsen, L., Bahr, R. Prevention of hamstring strains in elite soccer: an intervention study. *Scandinavian Journal of Medicine and Science in Sports*, 18: 40-48. 2008.
6. Askling, C., Karlsson, J. and Thorstensson, A. Hamstring injury occurrence in elite soccer players after preseason strength training with eccentric overload. *Scandinavian Journal of Medicine and Science in Sports*, 13: 244-250. 2003.
7. Baechle, T.R., Earle, R.W., and Wathen, D. Resistance training. In *Essentials of Strength Training and Conditioning*. T.R Baechle and R.W Earle Eds. Champaign, IL: Human Kinetics, 2008. pp. 381-412.
8. Bailey, R., Collins, D., Ford, P., MacNamara, A., Toms, M. and Pearce, G. Participant development in sport: An academic review. *Sports Coach UK*, 1-134. 2010.
9. Bailey, R. and Morley, D. Towards a model of talent development in physical education. *Sport, Education and Society*, 11: 211–230. 2006.
10. Baker, D., Mitchell, J., Boyle, D., Currell, S., Wilson, G., Bird, S.P., O'Connor, D. and Jones, J. Resistance training for children and youth: a position stand from the Australian Strength and Conditioning Association (ASCA). 2007. [www.strengthandconditioning.org](http://www.strengthandconditioning.org) (accessed 13 July 2011).
11. Balyi, I. and Hamilton, A. Long-Term Athlete Development: Trainability in Childhood and Adolescence - Windows of Opportunity - Optimal Trainability. Victoria: National Coaching Institute British Columbia & Advanced Training and Performance Ltd. 2004.

12. Barnett L., Van Beurden, E., Morgan, P., Brooks, L. and Beard, J. Childhood motor skill proficiency as a predictor of adolescent physical activity. *Journal of Adolescent Health*, 44: 252-259. 2009.
13. Bass, S.L. The prepubertal years – a unique opportune stage of growth when the skeleton is most responsive to exercise. *Sports Medicine*, 30: 73-78. 2000.
14. Bass SL and Myburg K. The effect of exercise on peak bone mass and bone strength. In *Sports Endocrinology*. M. Warren and N. Constantini Eds. Totowa, NJ: Humana Press Inc, 2000. pp. 253–280.
15. Bauer R and Steiner, M. Injuries in the European Union statistics summary 2005-2007. Vienna: European Network for Sports Injury Prevention and European Commission, Health and Consumers. 2009.
16. Baxter-Jones, A., Helms, P., Maffulli, N., Baines-Preece, J. and Preece, M. Growth and development of male gymnasts, swimmers, soccer and tennis Players: A longitudinal study. *Annals of Human Biology*, 22: 381-394. 1995.
17. Behm, D.G., Faigenbaum, A.D., Falk, B. and Klentrou, P. Canadian Society for Exercise Physiology position paper: resistance training in children and adolescents. *Applied Physiology Nutrition and Metabolism*, 33: 547-61. 2008.
18. Behringer, M., vom Heede, A., Matthews, M. and Mester, J. Effects of strength training on motor performance skills in children and adolescents: a meta-analysis. *Pediatric Exercise Science*, 23: 186-206. 2011.
19. Behringer, M., vom Heede, A., Yue, Z. and Mester, J. Effects of resistance training in children and adolescents: a meta-analysis. *Pediatrics*, 126: 1199-1210. 2010.
20. Beunen, G.P. Biological maturation and physical performance. In *Kinanthropometry IV*. W. Duquet and J.A.P. Day Eds. London: E & FN Spon, 1993. pp. 190-208.
21. Beunen, G.P. and Malina, R.M. Growth and biologic maturation: relevance to athletic performance. In *The Child and Adolescent Athlete*. H. Hebestreit and O. Bar-Or Eds. Oxford: Blackwell Publishing, 2005. pp. 3-17.
22. Beunen, G.P. and Malina, R.M. Growth and physical performance relative to the timing of the adolescent spurt. *Exercise and Sport Sciences Reviews*, 16: 503–540. 1988.
23. Beunen, G.P., Malina, R.M., Van't Hof, M.A., Simons, J., Ostyn, M., Renson, R. and Van Gerven, D. Adolescent Growth and Motor Performance. Champaign, IL: Human Kinetics, 1988. pp. 6–9.
24. Blimkie, C.J. Age- and sex-associated variation in strength during childhood: Anthropometric, morphologic, neurological, biomechanical, endocrinologic, genetic and physical activity correlates. In *Perspectives in Exercise Science and Sports*. C. Gisolfi and D. Lamb Eds. Indianapolis: Benchmark, 1989. pp. 99–163.
25. Blimkie, C.J., Rice, S., Webber, C.E., Martin, J., Levy, D. and Gordon, C.L. Effects of resistance training on bone mineral content and density in adolescent females. *Canadian Journal of Physiology and Pharmacology*, 74: 1025–1033. 1996.
26. Bloemers, F., Collard, D., Paw, M., Van Mechelen, W., Twisk, J. and Verhagen, E. Physical inactivity is a risk factor for physical activity-related injuries in children. *British Journal of Sports Medicine*. Epub ahead of print. 2011.
27. Bompa, T.O. *Total Training for Young Champions*. Champaign, IL, USA: Human Kinetics. 2000
28. Borms, J. The child and exercise: An overview. *Journal of Sports Sciences*, 4: 4–20. 1986.
29. Bouchant, A., Martin, V., Maffiuletti, N.A. and Ratel, S. Viewpoint: can muscle size fully account for strength differences between children and adults. *Journal of Applied Physiology*, 110: 1748-1749. 2011.
30. Branta, C., Haubenstricker, J. and Seefeldt, V. Age changes in motor skills during childhood and adolescence. *Exercise and Sport Sciences Reviews*, 12: 467-500. 1984.
31. Buchell, M., Mendez-Villanueva, A., Delhomel, G., Brughelli, M. and Ahmaidi, S. Improving sprint ability in young elite soccer players: repeated shuttle sprints vs. explosive strength training. *Journal of Strength and Conditioning Research* 24: 2715-2722. 2010.
32. Burgess, D. J. and Naughton, G. A. Talent development in adolescent team sports: a review. *International Journal of Sports Physiology and Performance*, 5: 103-116. 2010.
33. Byrd, R., Pierce, K., Reilly, L. and Brady, J. Young weightlifters' performance across time. *Sports Biomechanics*, 2: 133-140. 2003.
34. Cahill, B. and Griffith, E. Effect of preseason conditioning on the incidence and severity of high school football knee injuries. *American Journal of Sports Medicine*, 6: 180-184. 1978.
35. Casey, B.J., Tottenham, N., Liston, C. and Durston, S. Imaging the developing brain: what have we learned about cognitive development? *Trends in Cognitive Sciences*, 9: 104-110. 2005.
36. Casey BJ, Giedd JN and Thomas KM. Structural and functional brain development and its relation to cognitive development. *Biological Psychology*, 54: 241–257. 2000.
37. Chelly, M.S., Cherif, N., Amar, M.B., Hermassi, S., Fathloun, M., Bouhel, E., Tabka, Z. and Shepard, R.J. Relationships of peak leg power, 1 maximal repetition half back squat and leg muscle volume to 5-M sprint performance in junior soccer players. *Journal of Strength and Conditioning Research*, 24: 266-71. 2010.
38. Christou, M., Smilios, I., Sotiropoulos, K., Volaklis, K., Piliandis, T. and Tokmakidis, S.P. Effects of resistance training on the physical capacities of adolescent soccer players. *Journal of Strength and Conditioning Research*, 20: 783–791. 2006.
39. Clark, E., Tobias, J., Murray, L. and Boreham, C. Children with low muscle strength are at increased risk of fracture with exposure to exercise. *Journal of Musculoskeletal and Neuronal Interactions*, 11: 196-202. 2011.
40. Cohen, D.D., Voss, C., Taylor, M.J.D., Delextrat, A., Ogunleye, A.A. and Sandercock, G.R.H. Ten-year secular changes in muscular fitness in English children. *Acta Paediatrica*, 100: e175-177. 2011.
41. Cooper, R.M. and Zubek, J.P. Effects of enriched and restricted early environments on the learning ability of bright and dull rats. *Canadian Journal of Psychology*, 12: 159-164. 1958.
42. Conroy, B.P., Kraemer, W.J., Maresh, C.M., Fleck, S.J., Stone, M.H., Fry, A.C., Miller, P.D. and Dalsky, G.P. Bone mineral density in elite junior Olympic weightlifters. *Medicine and Science in Sports and Exercise*, 25: 1103-1109. 1993.
43. Cruz, M.L., Shaibi, G.Q., Weigensberg, M.J., Spruitz-Metz, D., Ball, G.D.C. and Goran, M.I. Pediatric obesity and insulin resistance: chronic disease risk and implications for treatment and prevention beyond body weight modification. *Annual Review of Nutrition*, 25: 435-468. 2005.
44. Davis, J.M., Tung, A., Chak, S.S., Ventura, E.E., Byrd-Williams, C.E., Alexander, K.E., Lane, C.J., Weigensberg, M.J., Spruijt-Metz, D. and Goran, M.I. Aerobic and strength training reduces adiposity in overweight Latina adolescents. *Medicine and Science in Sports and Exercise*, 41: 1494-1503. 2009.
45. De Ste Croix, M.B.A. Muscle strength. In *Paediatric exercise science and medicine*. N. Armstrong and W. Van Mechelen Eds. Oxford: Oxford University Press, 2008. pp. 199-211.
46. De Ste Croix, M.B.A. Advances in paediatric strength assessment: changing our perspective on strength development. *Journal of Sports Science and Medicine*, 6: 292-304. 2007.

47. D'hondt, E., Deforche, B., Vaeyens, R., Vandorpe, B., Vandendriessche, J., Pion, J., Philippaerts, R., de Bourdeaudhuij, I. and Lenoir, M. Gross motor coordination in relation to weight status and age in 5- to 12-year-old boys and girls: A cross sectional study. *International Journal of Pediatric Obesity*, 6: 556-564. 2011.
48. Dias Quiterio, A.L., Carnero, E.A., Baptista, F.M., and Sardinha, L.B. Skeletal mass in adolescent male athletes and nonathletes: relationships with high-impact sports. *Journal of Strength and Conditioning Research*, 25: 3439-3447. 2011.
49. DiStefano, L.J., Padua, D.A., Blackburn, J.T., Garrett, W.E., Guskiewicz, K.M., and Marshall, S.W. Integrated injury prevention program improves balance and vertical jump height in children. *Journal of Strength and Conditioning Research*, 24: 332-342. 2010.
50. Docherty, D., Wenger, H., Collis, M. and Quinney, H. The effects of variable speed resistance training on strength development in prepubertal boys. *Journal of Human Movement Studies*, 13: 377–382. 1987.
51. Dorgo, S., King, G.A., Candelaria, N.G., Bader, J.O., Brickey, G.D. and Adams, C.E. Effects of manual resistance training on fitness in adolescents. *Journal of Strength and Conditioning Research*, 23: 2287-2294. 2009.
52. Dvorkin, L.S. *Weightlifting and age: Scientific and Pedagogical Fundamentals of a Multi-Year System of Training Junior Weightlifters*. Translated by A. Charniga Jr. Livonia, MI: Sportivny Press, 1992.
53. Dvorkin, L.S. The Training of young weightlifters 13 -16 years old. In *The 1975 Russian Weightlifting Yearbook*. Translated by B.W Scheithauer. Moscow: Fiskultura I Sport Publishing, 1975. pp. 36-40.
54. Ekelund, U., Tomkinson, G. and Armstrong, N. What proportion of youth are physically active? Measurement issues, levels and recent time trends. *British Journal of Sports Medicine*. 45: 859-865. 2011.
55. Emery, C.A. and Meeuwisse, W. The effectiveness of a neuromuscular prevention strategy to reduce injuries in youth soccer: a cluster-randomised controlled trial. *British Journal of Sports Medicine*, 44: 555-562. 2010.
56. Eston, R., Byrne, C and Twist, C. Muscle function after exercise-induced muscle damage: Considerations for athletic performance in children and adults. *Journal of Exercise Science and Fitness*, 1: 85–96. 2003.
57. Faigenbaum A.D. Resistance training for children and adolescents: Are there health outcomes?. *American Journal of Lifestyle Medicine*, 1: 190-200. 2007.
58. Faigenbaum, A.D. Psychological benefits of prepubescent strength training. *Strength and Conditioning Journal*, 17: 28-32. 1995.
59. Faigenbaum, A.D. and Mediate, P. The effects of medicine ball training on physical fitness in high school physical education students. *The Physical Educator*, 63: 160-167. 2006.
60. Faigenbaum, A.D. and Myer, G.D. Resistance training among young athletes: Safety, efficacy and injury prevention effects. *British Journal of Sports Medicine*, 44: 56-63. 2010.
61. Faigenbaum, A.D. and Myer, G.D. Pediatric resistance training: benefits, concerns, and program design considerations. *Current Sports Medicine Reports*, 9: 161-168. 2010.
62. Faigenbaum, A.D. and Westcott, W.L. *Youth Strength Training*. Champaign, IL, USA: Human Kinetics. 2009.
63. Faigenbaum, A.D., Milliken, L.A. and Westcott, W.L. Maximal strength testing in healthy children. *Journal of Strength and Conditioning Research*, 17: 162-166. 2003.
64. Faigenbaum, A.D., Milliken, L., Moulton, L. and Westcott, W.L. Early muscular fitness adaptations in children in response to two different resistance training regimens. *Pediatric Exercise Science*, 17: 237–248. 2005.
65. Faigenbaum, A.D., Westcott, W.L., LaRousa Loud, R. and Long, C.J. The effects of different resistance training protocols on muscular strength and endurance development in children. *Pediatrics*, 104: e5. 1999.
66. Faigenbaum, A.D., Zaichkowsky, L., Westcott, W.L., Micheli, L.J. and Fehlandt, A. The effects of a twice per week strength training program on children. *Pediatric Exercise Science*, 5: 339–346. 1993.
67. Faigenbaum, A.D., LaRosa Loud, R., O'Connell, J., Glover, S., O'Connell, J. and Westcott, W.L. Effects of different resistance training protocols on upper-body strength and endurance development in children. *Journal of Strength and Conditioning Research*, 15: 459–465. 2001.
68. Faigenbaum, A.D., McFarland, J.E., Buchanan, E., Ratamess, N.A., Kang, J. and Hoffman, J.R. After-school fitness performance is not altered after physical education lessons in adolescent athletes. *Journal of Strength and Conditioning Research*, 24: 765–770. 2010.
69. Faigenbaum, A.D., Kraemer, W.J., Blimkie, C.J., Jeffreys, I., Micheli, L.J., Nitka, M. and Rowland, T.W. Youth resistance training: updated position statement paper from the National Strength and Conditioning Association. *Journal of Strength and Conditioning Research*, 23: S60-S79. 2009.
70. Faigenbaum, A.D., McFarland, J.E., Herman, R.E., Naclerio, F., Ratamess, N.A., Kang, J. and Myer, G.D. Reliability of the one-repetition-maximum power clean test in adolescent athletes. *Journal of Strength and Conditioning Research*, 26: 432-437. 2012.
71. Faigenbaum, A.D., McFarland, J., Keiper, F., Tevlin, W., Kang, J., Ratamess, N. and Hoffman, J. Effects of a short term plyometric and resistance training program on fitness performance in boys age 12 to 15 years. *Journal of Sports Science and Medicine*, 6: 519–525. 2007.
72. Faigenbaum, A.D., Ratamess, N., McFarland, J., Kaczmarek, J., Coraggio, M., Kang, J. and Hoffman, J. Effect of rest interval length on bench press performance in boys, teens and men. *Pediatric Exercise Science*, 20: 457–469. 2008.
73. Faigenbaum, A.D., Westcott, W.L., Micheli, L.J., Outerbridge, R.A., Long, C.J., LaRosa-Loud, R. and Zaichkowsky, L.D. The effects of strength training and detraining on children. *Journal of Strength and Conditioning Research*, 10: 109-14. 1996.
74. Falk, B. and Dotan, R. Child-adult differences in the recovery from high intensity exercise. *Exercise and Sport Sciences Reviews*, 34: 107–112. 2006.
75. Falk, B. and Eliakim, A. Resistance training, skeletal muscle and growth. *Paediatric Endocrinology Reviews*, 1: 120-27. 2003.
76. Falk, B. and Tenenbaum, G. The effectiveness of resistance training in children: a meta-analysis. *Sports Medicine*, 22: 176-186. 1996.
77. Ford, K.R., Myer, G.D. and Hewett, T.E. Longitudinally decreased knee abduction and increased hamstrings strength in females with self-reported resistance training. *Proceedings of the American College of Sports Medicine Annual Meeting*. Denver, Colorado. 2011.
78. Ford, K.R., Shapiro, R., Myer, G.D., Van Den Bogert, A.J. and Hewett, T.E. Longitudinal Sex Differences during Landing in Knee Abduction in Young Athletes. *Medicine and Science in Sports and Exercise*, 42: 1923-1931. 2010.
79. Ford, K.R., Myer, G.D., Smith, R.L., Byrnes, R.N., Dopirak, S.E. and Hewett, T.E. Use of an overhead goal alters vertical jump performance and biomechanics. *Journal of Strength and Conditioning Research*, 19: 394-399. 2005.
80. Ford, P.A., De Ste Croix, M.B.A., Lloyd, R.S., Meyers, R., Moosavi, M., Oliver, J., Till, K. and Williams, C.A. The Long-Term Athlete Development model: Physiological

- evidence and application. *Journal of Sports Science*, 29: 389-402. 2011.
81. Fuchs, R.K., Bauer, J.J. and Snow, C.M. Jumping improves hip and lumbar spine bone mass in prepubescent children: A randomized controlled trial. *Journal of Bone Mineral Research*, 16: 148–156. 2001.
  82. Fukunaga T., Funato K. and Ikegawa S. The effects of resistance training on muscle area and strength in prepubertal age. *Annals of Physiology and Anthropology* 11: 357-64, 1992.
  83. Gabbe, B.J., Branson, R. and Bennell, K.L. A pilot randomised controlled trial of eccentric exercise to prevent hamstring injuries in community-level Australian Football. *Journal of Science and Medicine in Sport*, 9: 103-109. 2006.
  84. Gortmaker, S., Lee, R., Craddock, A., Sobol, A., Duncan, D. and Wang, Y. Disparities in Youth Physical Activity in the United States: 2003-2006. *Medicine and Science in Sports and Exercise*, epub ahead of print. 2011.
  85. Granacher, U., Goeseles, A., Roggo, K., Wischer, T., Fischer, S., Zuerny, C., Gollhofers, A. and Kriemler, S. Effects and mechanisms of strength training in children. *International Journal of Sports Medicine*, 32: 357-364. 2011a.
  86. Granacher, U., Muehlbauer, T., Doerflinger, B., Strohmeier, R., and Gollhofer, A. Promoting strength and balance in adolescents during physical education: effects of a short-term resistance training. *Journal of Strength and Conditioning Research*, 25: 940-949. 2011b.
  87. Gunter, K., Almstedt, H. and Janz, K. Physical activity in childhood may be the key to optimizing lifespan skeletal health. *Exercise and Sports Science Reviews*, 40: 13-21. 2011.
  88. Guthold, R., Cowan, M., Autenrieth, C., Kann, L. and Riley, L. Physical activity and sedentary behavior among schoolchildren: a 34 country comparison. *Journal of Pediatrics*, 157: 43-49. 2010.
  89. Häkkinen, K., Mero, A. and Kauhanen, H. Specificity of endurance, sprint and strength training on physical performance capacity in young athletes. *Journal of Sports Medicine*, 29: 27-35. 1989.
  90. Hamill, B. Relative safety of weightlifting and weight training. *Journal of Strength and Conditioning Research*, 8: 53-57. 1994.
  91. Hands B. Changes in motor skill and fitness measures among children with high and low motor competence: a five-year longitudinal study. *Journal of Science and Medicine in Sport*, 11: 155-162. 2008.
  92. Hass C.J., Feigenbaum, M.S and Franklin, B.A. Prescription of resistance training for healthy populations. *Sports Medicine*, 31: 953-64. 2001.
  93. Hejna, W.F., Rosenberg, A., Buturusis, D.J. and Krieger, A. The prevention of sports injuries in high school students through strength training. *National Strength Coaches Association Journal*, 4: 28-31. 1982.
  94. Hetherington, M. Effect of isometric training on the elbow flexion force torque of grade five boys. *Research Quarterly*, 47: 41–47. 1976.
  95. Hewett, T.E., Ford, K.R. and Myer, G.D. Anterior Cruciate Ligament Injuries in Female Athletes: Part 2, A Meta-analysis of Neuromuscular Interventions Aimed at Injury Prevention. *American Journal of Sports Medicine*, 34: 490-498. 2006.
  96. Hewett, T.E., Myer, G.D. and Ford, K.R. Reducing knee and anterior cruciate ligament injuries among female athletes: a systematic review of neuromuscular training interventions. *The Journal of Knee Surgery*, 18: 82-88. 2005.
  97. Hewett TE, Myer GD, Ford KR. Decrease in neuromuscular control about the knee with maturation in female athletes. *Journal of Bone and Joint Surgery*, 86: 1601-1608. 2004.
  98. Hewett, T.E., Lindenfeld, T.N., Riccobene, J.V. and Noyes, F.R. The effect of neuromuscular training on the incidence of knee injury in female athletes. A prospective study. *American Journal of Sports Medicine*, 27: 699-706. 1999.
  99. Hewett, T.E., Myer, G.D., Ford, K.R. and Slauterbeck, J.R. Preparticipation Physical Exam Using a Box Drop Vertical Jump Test in Young Athletes: The Effects of Puberty and Sex. *Clinical Journal of Sports Medicine*, 16: 298-304. 2006.
  100. Hewett, T.E., Stroupe, A.L, Nance, T.A. and Noyes, F.R. Plyometric training in female athletes. Decreased impact forces and increased hamstring torques. *American Journal of Sports Medicine*, 24: 765-773. 1996.
  101. Hewett, T.E., Myer, G.D., Ford, K.R., Heidt, R.S., Colosimo, A.J., McLean, S.G., van den Bogert, A.J., Paterno, M.V. and Succop, P. Biomechanical Measures of Neuromuscular Control and Valgus Loading of the Knee Predict Anterior Cruciate Ligament Injury Risk in Female Athletes: A Prospective Study. *American Journal of Sports Medicine*, 33: 492-501. 2005.
  102. Hind, K. and Burrows, M. Weight-bearing exercise and bone mineral accrual in children and adolescents: a review of controlled trials. *Bone*, 40: 14-27. 2007.
  103. Holloway, J., Beuter, A. and Duda, J. Self-efficacy and training in adolescent girls. *Journal of Applied Social Psychology*, 18: 699–719. 1988.
  104. Horvat, M., Franklin, C. and Born, D. Predicting strength in high school women athletes. *Journal of Strength and Conditioning Research*, 21: 1018–1022. 2007.
  105. Ingle, L., Sleap, M. and Tolfrey, K. The effect of a complex training and detraining programme on selected strength and power variables in early pubertal boys. *Journal of Sports Sciences*, 24: 987–997. 2006.
  106. Jones, R.M., Fry, A.C., Weiss, L.W., Kinzey, S.J. and Moore, C.A. Kinetic comparison of free weight and machine power cleans. *Journal of Strength and Conditioning Research*, 22: 1785–1789. 2008.
  107. Jonhagen, S., Nemeth, G. and Eriksson, E. Hamstring injuries in sprinters: the role of concentric and eccentric hamstring muscle strength and flexibility. *American Journal of Sports Medicine*, 22: 262-266. 1994.
  108. Kaufman, L.B. and Schilling, D.L. Implementation of a strength training program for a 5-year-old child with poor body awareness and developmental coordination disorder. *Physical Therapy*, 87: 455-467. 2007.
  109. Kawamori, N. and Newton, R.U. Velocity specificity of resistance training: actual movement velocity versus intention to move explosively. *Strength and Conditioning Journal*, 28: 86-91. 2006.
  110. Keiner, M., Sander, A., Wirth, K., Caruso, O., Immesberger, P. and Zawieja, M. Trainability of children and adolescents in the front and back squat. *Journal of Strength and Conditioning Research*, in press.
  111. Kirk, D. Physical education, youth sport and lifelong participation: the importance of early learning experiences. *European Physical Education Review*, 11: 239-255. 2005.
  112. Kraemer, W.J., Fry, A.C., Frykman, P.N., Conroy, B. and Hoffman, J. Resistance training and youth. *Pediatric Exercise Science*, 1: 336-350. 1989.
  113. Kraemer, W.J., Keuning, M., Ratamess, N.A., Volek, J.S., McCormick, M., Bush, J.A., Nindi, B.C., Gordon, S.E., Mazzetti, S.A., Newton, R.U., Gómez, A.L., Wickham, R.B., Rubin, M.R. and Häkkinen, K. Resistance training combined with bench-step aerobics enhances women's health profile. *Medicine and Science in Sports and Exercise*, 33: 259-269. 2001.
  114. Kravitz, L., Akalan, C., Nowicki, K. and Kinzey, S. Prediction of 1 repetition maximum in high school power lifters. *Journal of Strength and Conditioning Research*, 17:

- 167–172. 2003.
115. Lau, P.W.C., Kong, Z., Choi, C., Yu, C.C.W., Chan, D.F.Y., Sung, R.Y.T. and Leung, B.W.C. Effects of short-term resistance training on serum leptin levels in obese adolescents. *Journal of Exercise Science and Fitness*, 8: 54-60. 2010.
  116. Lehnhard, R.A., Lehnhard, H.R., Young, R. and Butterfield, S.A. Monitoring injuries on a college soccer team: the effect of strength training. *Journal of Strength and Conditioning Research*, 10: 115-119. 1996.
  117. Lephart, S.M., Abt, J.P., Ferris, C.M., Sell, C.T., Nagai, T., Meyers, J.B. and Irrgang, J.J. Neuromuscular and biomechanical characteristic changes in high school athletes: a plyometric versus basic resistance program. *British Journal of Sports Medicine*, 39: 932-938. 2005.
  118. Lillegard, W.A., Brown, E.W., Wilson, D.J., Henderson, R. and Lewis, E. Efficacy of strength training in prepubescent to early postpubescent males and females: effects of gender and maturity. *Pediatric Rehabilitation*, 1: 147-157. 1997.
  119. Lloyd, R.S. and Oliver, J.L. The Youth Physical Development model: a new approach to long-term athletic development. *Strength and Conditioning Journal* 34: 37-43. 2012.
  120. Lloyd, R.S., Oliver, J.L., Hughes, M.G. and Williams, C.A. Effects of 4-weeks plyometric training on reactive strength index and leg stiffness in male youths. *Journal of Strength and Conditioning Research*, in press.
  121. Lloyd, R.S., Oliver, J.L., Meyers, R.W., Moody, J. and Stone, M.H. Long-term athletic development and its application to youth weightlifting. *Strength and Conditioning Journal*, in press.
  122. Lopes, V., Rodrigues, L., Maia, J. and Malina, R. Motor coordination as predictor of physical activity in childhood. *Scandinavian Journal of Medicine and Science in Sports*, 21: 663-669. 2011.
  123. Lopes, V., Stodden, D., Bianchi, M., Maia, J. and Rodrigues, L. Correlation between BMI and motor coordination in children. *Journal of Science and Medicine in Sport*, 15: 38-43. 2012.
  124. Lubans, D.R., Morgan, P.J., Cliff, D.P., Barnett, L.M. and Okely, A.D. Fundamental movement skills in children and adolescents. *Sports Medicine*, 40: 1019–1035. 2010.
  125. Malina, R. Weight training in youth-growth, maturation, and safety: an evidence-based review. *Clinical Journal of Sports Medicine*, 16: 478-487. 2006.
  126. Malina, R.M., Bouchard, C. and Bar-Or, O. *Growth, Maturation, and Physical Activity*. Champaign, IL, USA: Human Kinetics. 2004 pp. 3-20.
  127. Matos, N., and Winsley, R.J. Trainability of young athletes and overtraining. *Journal of Sports Science and Medicine*, 6: 353-367. 2007.
  128. Mayhew, J., Kerksick, C., Lentz, D., Ware, J. and Mayhew, D. Using repetitions to predict one-repetition maximum bench press in male high school athletes. *Pediatric Exercise Science*, 16: 265–276. 2004.
  129. McGuigan, M.R., Tatasciore, M., Newton, R.U. and Pettigrew, S. Eight weeks of resistance training can significantly alter body composition in children who are overweight or obese. *Journal of Strength and Conditioning Research*, 23: 80-85. 2009.
  130. McHugh, M. Oversized young athletes: a weighty concern. *British Journal of Sports Medicine*, 44: 45-49. 2010.
  131. Mersch, F. and Stoboy, H. Strength training and muscle hypertrophy in children. In *Children and Exercise XIII*. S. Oseid and K. Carlsen Eds. Champaign, IL: Human Kinetics Books, 1989. pp. 165–182.
  132. Meylan, C. and Malatesta, D. Effects of in-season plyometric training within soccer practice on explosive actions of young players. *Journal of Strength and Conditioning Research*, 23: 2605-2613. 2009.
  133. Micheli, L. Preventing injuries in team sports: What the team physician needs to know. In *F.I.M.S. Team Physician Manual (2nd edition)*. K. Chan, L. Micheli, A. Smith, C. Rolf, N. Bachl, W. Frontera and T. Alenabi Eds. Hong Kong: CD Concepts, 2006. pp. 555-572.
  134. Mikkola, J., Rusko, H., Nummela, A., Pollari, T. and Häkkinen, K. Concurrent endurance and explosive type strength training improves neuromuscular and anaerobic characteristics in young distance runners. *International Journal of Sports Medicine*, 28: 602-611. 2007.
  135. Moliner-Urdiales, D., Ruiz, J.R., Ortega, F.B., Jiménez-Pavón, D., Vicente-Rodriguez, G., Rey-López, J.P., Martínez-Gómez, D., Casajús, J.A., Mesana, M.I., Marcos, A., Noriega-Borge, M.J., Sjöström, M., Castillo, M.J. and Moreno, L.A. Secular trends in health-related physical fitness in Spanish adolescents: the AVENA and HELENA studies. *Journal of Science and Medicine in Sport*, 13: 584-588. 2010.
  136. Mountjoy, M., Andersen, L., Armstrong, N., Biddle, S., Boreham, C., Bedenbeck, H.P.B., Ekelund, U., Engbretsen, L., Hardman, K., Hills, A., Kahlmeier, S., Kriemler, S., Lambert, E., Ljungqvist, A., Matsudo, V., McKay, H., Micheli, L., Pate, R., Riddoch, C., Schamasch, P., Sundberg, C.J., Tomkinson, G., van Sluijs, E. and van Mechelen, W. International Olympic Committee Consensus statement on the health and fitness of young people through physical activity and sport. *British Journal of Sports Medicine*, 45:839-848. 2011.
  137. Myer, G.D., Sugimoto, D. and Hewett, T.E. The influence of age on the effectiveness of neuromuscular training to reduce anterior cruciate ligament injury in female athletes: A Meta-Analysis. Accepted For Presentation at the National Athletic Trainer's Association (NATA) National Meeting. 2012.
  138. Myer, G.D., Ford, K.R., Brent, J.L. and Hewett, T.E. Differential neuromuscular training effects on ACL injury risk factors in "high-risk" versus "low-risk" athletes. *BMC Musculoskeletal Disorders*, 8: 1-7. 2007.
  139. Myer, G.D., Ford, K.R., Brent, J.L. and Hewett, T.E. The effects of plyometric versus dynamic balance training on power, balance and landing force in female athletes. *Journal of Strength and Conditioning Research*, 20: 345-353. 2006.
  140. Myer, G.D., Ford, K.R., McLean, S.G. and Hewett, T.E. The effects of plyometric versus dynamic stabilization and balance training on lower extremity biomechanics. *American Journal of Sports Medicine*, 34: 490-498. 2006.
  141. Myer, G.D., Ford, K.R., Palumbo, J.P. and Hewett, T.E. Neuromuscular training improves performance and lower-extremity biomechanics in female athletes. *Journal of Strength and Conditioning Research*, 19: 51-60. 2005.
  142. Myer, G.D., Quatman, C.E., Khoury, J., Wall, E.J. and Hewett, T.E. Youth versus adult "weightlifting" injuries presenting to United States emergency rooms: accidental versus nonaccidental injury mechanisms. *Journal of Strength and Conditioning Research*, 23: 2054-2060. 2009.
  143. Myer, G.D., Faigenbaum, A.D., Chu, D., Falkel, J., Ford, K. and Best, T. Integrative training for children and adolescents: techniques and practices for reducing sports-related injuries and enhancing athletic performance. *The Physician and Sportsmedicine*, 39: 74-84. 2011
  144. Myer, G.D., Faigenbaum, A.D., Ford, K.R., Best, T.M., Bergeron, M.F. and Hewett, T.E. When to initiate integrative neuromuscular training to reduce sport-related injuries and enhance health in youth. *Current Sports Medicine Reports*, 10: 157-166. 2011.
  145. Myer, G.D., Ford, K.R., Barber Foss, K.D., Liu, C., Nick, T.G. and Hewett, T.E. The relationship of hamstrings and quadriceps strength to anterior cruciate ligament injury in female athletes. *Clinical Journal of Sports Medicine*, 19: 3-

8. 2009.
146. Myer, G.D., Ford, K.R., Barber Foss, K.D., Goodman, A., Ceasar, A., Rauh, M.J. Divine, J.G. and Hewett, T.E. The incidence and potential pathomechanics of patellofemoral pain in female athletes. *Clinical Biomechanics*, 25: 700-707. 2010.
147. Myklebust, G., Engebretsen, L., Braekken, I.H., Skjølberg, A., Olsen, O.E. and Bahr, R. Prevention of anterior cruciate ligament injuries in female team handball players: a prospective intervention study over three seasons. *Clinical Journal of Sports Medicine*, 13: 71-78. 2003.
148. Nader, P., Bradley, R., Houts, R., McRitchie, S. and O'Brien, M. Moderate to vigorous physical activity from ages 9 to 15 years. *The Journal of the American Medical Association*, 300: 295-305. 2008.
149. Naughton, G., Farpour, L., Carlson, J., Bradney, M. and Van Praagh, E. Physiological issues surrounding the performance of adolescent athletes. *Sports Medicine*, 30: 309-325. 2000.
150. Naylor, L.H., Watts, K., Sharpe, J.A., Jones, T.W., Davis, E.A., Thompson, A., George, K., Ramsay, J.M., O'Driscoll, G. and Green, D.J. Resistance training and diastolic myocardial tissue velocities in obese children. *Medicine and Science in Sports and Exercise*, 40: 2027-2032. 2008.
151. Neu, C.M., Rauch, F., Rittweger, J., Manz, F. and Schoenau, E. Influence of puberty on muscle development at the forearm. *American Journal of Physiology – Endocrinology and Metabolism*, 283: E103 – E107. 2002.
152. Nichols, D.L., Snaborn, C.F. and Love, A.M. Resistance training and bone mineral density in adolescent females. *Journal of Pediatrics*, 139: 494-500. 2001.
153. Norris, S.R. Long-term athlete development Canada: attempting system change and multi-agency cooperation. *Current Sports Medicine Reports*, 9: 379-82. 2010.
154. Nyberg, G., Nordenfelt, A., Ekelund, U. and Marcus, C. Physical activity patterns measured by accelerometry in 6- to 10-yr-old children. *Medicine and Science in Sports and Exercise*, 41: 1842-1848. 2009.
155. O'Brien, T.D., Reeves, N.D., Baltzopoulos, V., Jones, D.A. and Magnaris, C.N. In vivo measurements of muscle specific tension in adults and children. *Experimental Physiology*, 95: 202-210. 2010.
156. O'Brien, T.D., Reeves, N.D., Baltzopoulos, V., Jones, D.A. and Magnaris, C.N. Strong relationships exist between muscle volume, joint power and whole-body external mechanical power in adults and children. *Experimental Physiology*, 94: 731-738. 2009.
157. Orchard, J., Marsden, J., Lord, S. and Garlick, D. Preseason hamstring muscle weakness associated with hamstring muscle injury in Australian footballers. *American Journal of Sports Medicine*, 25: 81-85. 1997.
158. Ortega, F., Ruiz, J., Castillo, M. and Sjostrom, M. Physical fitness in children and adolescence: a powerful marker of health. *International Journal of Obesity*, 32: 1-11. 2008.
159. Ozmun, J.C., Mikesky, A.E. and Surburg, P. Neuromuscular adaptations following prepubescent strength training. *Medicine and Science in Sports and Exercise*, 26: 510-514. 1994.
160. Padilla-Moledo, C., Ruiz, J.R., Ortega, F.B., Mora, J. and Castro-Pinero, J. Associations of muscular fitness with psychological positive health, health complaints, and health risk behaviors in Spanish children and adolescents. *Journal of Strength and Conditioning Research*, 26: 1671-1673. 2012.
161. Parker, D.F., Round, J.M., Sacco, P. and Jones, D.A. A cross-sectional survey of upper and lower limb strength in boys and girls during childhood and adolescence. *Annals of Human Biology*, 17: 199-211. 1990.
162. Payne, V.G., Morrow, J.R., Johnson, L. and Dalton, S.N. Resistance training in children and youth. *Research Quarterly*, 68: 80-88. 1997.
163. Petersen, J., Thorborg, K., Nielsen, M.B., Budtz-Jorgensen, E. and Holmich, P. Preventive effect of eccentric training on acute hamstring injuries in men's soccer: a cluster-randomized controlled trial. *American Journal of Sports Medicine*, 39: 2296-2303. 2011.
164. Peterson M.D., Rhea M.R., Alvar, B.A. Applications of the dose-response for muscular strength development: a review of meta-analytic efficacy and reliability for designing training prescription. *Journal of Strength and Conditioning Research*, 19: 950-958. 2005.
165. Pfeiffer, R. and Francis, R. Effects of strength training on muscle development in prepubescent, pubescent and postpubescent males. *The Physician and Sports Medicine*, 14: 134-143. 1986.
166. Pierce, K.C., Byrd, R. and Stone, M.H. Youth weightlifting – is it safe?. *Weightlifting USA*, 17: 5. 1999.
167. Pierce, K.C., Brewer C., Ramsey, M.W., Byrd, R., Sands, W.A., Stone, M.E. and Stone, M.H. Youth resistance Training. *Professional Strength and Conditioning Journal*, 10: 9-23. 2008.
168. Quatman, C.E., Ford, K.R., Myer, G.D. and Hewett, T.E. Maturation leads to gender differences in landing force and vertical jump performance: a longitudinal study. *American Journal of Sports Medicine*, 34: 806-813. 2006.
169. Rabinowicz, T. The differentiated maturation of the cerebral cortex. In *Human Growth: A Comprehensive Treatise, Postnatal Growth: Neurobiology*. F. Falkner and J. Tanner Eds (volume 2). New York, NY: Plenum. 1986.
170. Ramsay JA, Blimkie CJR, Smith K, Garner S, MacDougall JD, Sale DG. Strength training effects in prepubescent boys. *Medicine and Science in Sports and Exercise*, 22: 605-614. 1990.
171. Rogasch, N.C., Dartnall, T.J., Cirillo, J., Nordstrom, M.A. and Semmler, J.G. Corticomotor plasticity and learning of a ballistic thumb training task are diminished in older adults. *Journal of Applied Physiology*, 107: 1874-1883. 2009.
172. Rosengren, K.S., Geert, J.P. and Savelsbergh, J.K. Development and learning: a TASC-based perspective of the acquisition of perceptual-motor behaviors. *Infant Behavior and Development*, 26: 473-494. 2003.
173. Runhaar, J., Collard, D.C.M., Kemper, H.C.G., van Mechelen, W. and Chinapaw, M. Motor fitness in Dutch youth: differences over a 26-year period (1980-2006). *Journal of Science and Medicine in Sport*, 13: 323-328. 2010.
174. Sadres, E., Eliakim, A., Constantini, N., Lidor, R. and Falk, B. The effect of long-term resistance training on anthropometric measures, muscle strength, and self concept in pre-pubertal boys. *Pediatric Exercise Science*, 13: 357-372. 2001.
175. Schick, E.E., Coburn, J.W., Brown, L.E., Judelson, D.A., Khamoui, A.V., Tran, T.T. and Uribe, B.P. A comparison of muscle activation between a Smith machine and free weight bench press. *Journal of Strength and Conditioning Research*, 24: 779-784. 2010.
176. Schwanbeck, S., Chilibeck, P.D. and Binsted, G. A comparison of free weight squat to smith machine squat using electromyography. *Journal of Strength and Conditioning Research*, 23: 2588-2591. 2009.
177. Schwingshandl, J., Sudi, K., Eibl, B., Wallner, S. and Borkenstein, M. Effect of individualised training programme during weight reduction on body composition: a randomised trial. *Archives of Disease in Childhood*, 81: 426-428. 1999.
178. Sgro, M., McGuigan, M.R., Pettigrew, S. and Newton, R.U. The effect of duration of resistance training interventions in children who are overweight or obese. *Journal of Strength and Conditioning Research*, 23: 1263-1270. 2009.
179. Shaibi, G., Cruz, M., Ball, G., Weigensberg, M., Salem, G.,

- Crespo, N. and Goran, M. Effects of resistance training on insulin sensitivity in overweight Latino adolescent males. *Medicine and Science in Sports and Exercise*, 38: 1208–1215. 2006.
180. Soligard, T., Myklebust, G., Steffen, K., Holme, I., Silvers, H., Bizzini, M., Junge, A., Dvorak, J., Bahr, R. and Andersen, T.E. Comprehensive warm-up programme to prevent injuries in young female footballers: cluster randomized controlled trial. *British Medical Journal*, 337: a2469. 2008.
181. Sothorn, M.S., Loftin, M.J., Udall, J.N., Suskind, R.M., Ewing, T.L., Tang, S.C. and Blecker, U. Safety, feasibility, and efficacy of a resistance training program in preadolescent obese children. *American Journal of the Medical Sciences*, 319: 370-375. 2000.
182. Stodden, D., Langendorfer, S. and Robertson, M. The association between motor skill competence and physical fitness in young adults. *Research Quarterly for Exercise and Sport*, 80: 223-229. 2009.
183. Stone, M.H., Pierce, K.C., Sands, W.A. and Stone, M.E. Weightlifting: a brief overview. *Strength and Conditioning Journal*, 28: 50-66. 2006.
184. Stratton G and Williams CA. Children and Fitness Testing. In *Sport and Exercise Physiology Testing Guidelines: the British Association of Sport and Exercise Sciences Guide*. E.M. Winter, A.M. Jones, R.C.R. Davison, P.D. Bromley and T.H. Mercer Eds. Oxon: Routledge, 2007. pp. 211-223.
185. Stratton, G., Jones, M., Fox, K.R., Tolfrey, K., Harris, J., Maffulli, N., Lee, M. and Frostick, S.P. Position Statement on guidelines for resistance exercise in young people. *Journal of Sports Science*, 22: 383-90. 2004.
186. Suh, S., Jeong, I.K., Kim, M.Y., Kim, Y.S., Shin, S., Kim, S.S. and Kim, J.H. Effects of resistance training and aerobic exercise on insulin sensitivity in overweight Korean adolescents: a controlled randomized trial. *Diabetes and Metabolism Journal*, 35: 418-426. 2011.
187. Szymanski, D.J., Szymanski, J.M., Molloy, J.M. and Pascoe, D.D. Effect of 12 weeks of wrist and forearm training on high school baseball players. *Journal of Strength and Conditioning Research*, 18: 432– 440. 2004.
188. Thomas, K., French, D. and Hayes, P.R. The effect of plyometric training techniques on muscular power and agility in youth soccer players. *Journal of Strength and Conditioning Research*, 23: 332–335. 2009.
189. Tonson, A., Ratel, S., Le Fur, Y., Cozzone, P. and Bendahan, D. Effect of maturation on the relationship between muscle size and force production. *Medicine and Science in Sports and Exercise*, 40: 918-925. 2008.
190. Tsolakis, C.K., Vagenas, G.K. and Dessypris, A.G. Strength adaptations and hormonal responses to resistance training and detraining in preadolescent males. *Journal of Strength and Conditioning Research*, 18: 625– 629. 2004.
191. Tsolakis, C., Messinis, D., Stergioulas, A. and Dessypris, A. Hormonal responses after strength training and detraining in prepubertal and pubertal boys. *Journal of Strength and Conditioning Research*, 14: 399-404. 2000.
192. Tudor-Locke, C., Johnson, W.D. and Katzmarzyk, P.T. Accelerometer-determined steps per day in US children and youth. *Medicine and Science in Sports and Exercise*, 42: 2244-2250. 2010.
193. Valovich-McLeod, T.C., Decoster, L.C., Loud, K.J., Micheli, L., Parker, J.T., Sandrey, M.A. and White C. National Athletic Trainers' Association position statement: prevention of pediatric overuse injuries. *Journal of Athletic Training*, 46: 206-220. 2011.
194. Van der Heijden, G., Wang, Z., Chu, Z., Toffolo, G., Manesso, E., Sauer, P.J. and Sunehag, A.L. Strength exercise improves muscle mass and hepatic insulin sensitivity in obese youth. *Medicine and Science in Sports and Exercise*, 42: 1973-1980. 2010.
195. Velez, A., Golem, D.L. and Arent, S.M. The impact of a 12-week resistance training program on strength, body composition, and self-concept of hispanic adolescents. *Journal of Strength and Conditioning Research*, 24: 1065-1073. 2010.
196. Verschuren, O., Ada, L., Maltais, D.B., Gorter, J.W., Scianni, A. and Ketelaar, M. Muscle strengthening in children with spastic cerebral palsy: considerations for future resistance training protocols. *Physical Therapy*, 91: 1130-1139. 2011.
197. Vicente-Rodriguez, G. How does exercise affect bone development during growth? *Sports Medicine*. 36: 561-569. 2006.
198. Viru A., Loko, J., Harro, M., Volver, A., Laaneots, L. and Viru, M. Critical periods in the development of performance capacity during childhood and adolescence. *European Journal of Physical Education*, 4: 75–119. 1999.
199. Virvidakis, K., Georgiu, E., Korkotsidis, A., Ntalles, A. and Proukakis, C. Bone mineral content of junior competitive weightlifters. *International Journal of Sports Medicine*, 11: 244-246. 1990.
200. Watts, K., Beye, P., Siafarikas, A., Davis, E., Jones, T., O'Driscoll, G. and Green, D. Exercise training normalizes vascular dysfunction and improves central adiposity in obese adolescents. *Journal of the American College of Cardiology*, 43: 1823–1827. 2004.
201. Weltman, A., Janney, C., Rians, C., Strand, K., Berg, B., Tippit, S., Wise, J., Cahill, B. and Katch, F. The effects of hydraulic resistance strength training in pre-pubertal males. *Medicine and Science in Sports and Exercise*, 18: 629–638. 1986.
202. Witzke, K.A. and Snow, C.M. Effects of plyometric jump training on bone mass in adolescent girls. *Medicine and Science in Sports and Exercise*, 32: 1051–1057. 2000.
203. World Health Organization. *Global Recommendations on Physical Activity for Health*. Geneva: WHO Press. 2010.
204. Yamamoto, T. Relationship between hamstring strains and leg muscle strength. A follow-up study of collegiate track and field athletes. *Journal of Sports Medicine and Physical Fitness*, 33: 194-199. 1993.
205. Young, W.B. Transfer of strength and power training to sports performance. *International Journal of Sports Physiology and Performance*, 1: 74-83. 2006.
206. Yu, C., Sung, R., Hau, K., Lam, P., Nelson, E. and So, R. The effect of diet and strength training on obese children's physical self concept. *Journal of Sports Medicine and Physical Fitness*, 48: 76–82. 2008.
207. Yu, C.C.W., Sung, R.Y.T., So, R.C.H., Lui, K.C., Lau, W., Lam, P.K.W. and Lau, E.M.C. Effects of strength training on body composition and bone mineral content in children who are obese. *Journal of Strength and Conditioning Research*, 19: 667-672. 2005.
208. Zafeiridis, A., Dalamitros, A., Dipla, K., Manou, V., Galanis, N. and Kellis, S. Recovery during high-intensity intermittent anaerobic exercise in boys, teens and men. *Medicine and Science in Sports and Exercise*, 37: 505–512. 2005.
209. Zakas A., Doganis, G., Papageorgopoulou, M., Zakas, N. and Vamvakoudis, E. The effect of cycle ergometer strength training in pubescent and post-pubescent untrained males. *Isokinetics and Exercise Science*, 14: 45–52. 2004.